geodesy.noaa.gov

Towards a More Consistent Geoid Model for North America (9619)

Kevin M. Ahlgren¹, Yan-Ming Wang¹, Xiaopeng Li², Monica A. Youngman¹ (USA)

¹NOAA's National Geodetic Survey ²ERT Inc.



Additional Geoids in US Pacific Islands



International Cooperation & Collaboration

- National Geospatial-Intelligence Agency (NGA)
 - NGA Surface Gravity Data where allowable
 - NGS GRAV-D data
 - NGA Preliminary Geopotential Models (PGMs)
- Geodetic groups within North America
 - Canadian Geodetic Survey (CGS) & Mexico's Instituto Nacional de Estadística y Geografía (INEGI)
- Other Groups
 - Agreements for data from Greenland (DTU), Iceland, etc.

geodesy.noaa.gov

NGA: Surface Gravity Data Sharing



geodesy.noaa.gov

NGA: Surface Gravity Data Sharing



NGA: GRAV-D Sharing

- GRAV-D major source of new data for NGA's EGM2020
- NGS will share 'cleaned' data annually (~June)
- Fits into NGA reference model schedule for EGM2020.



Data Blocks Ready for PGM18 (green)

NGA: PGM Validation

- EGM2020 release in Dec. 2019
- Annual PGM
 reference models
- PGM17 released in Dec. 2017
- Nearly identical to NGS REF17B model



Common Datasets: CGS, INEGI, & NGS

- Differences in North American Geoid Models:
 - 1) Different modeling (theory)
 - 2) Different input datasets
- Project initiated to address 2) \rightarrow Common Datasets
 - Surface gravity data (terrestrial, shiptrack, altimetry)
 - Airborne gravity data (GRAV-D)
 - DEM
 - Satellite gravity model

Common Datasets: Terrestrial Gravity

- Each group provide data over their territory
- v1 Completed
- v2 to include supplemental areas (Iceland, South America, Greenland, etc.)
- INEGI: Central & South America
- CGS: Greenland
- NGS: Carribbean



Common Datasets: DEM

- 3" elevation grid
- Mainly consists of SRTMv4.1 (below 60° N).
- Supplemental grids in other areas
- v1 completed
- v2 completed over the next year to include data over Alaska, Greenland, etc.



Common Datasets: DEM – Void Fills

- SRTMv4.1 still has areas that are not correct
 - Shadows incorrectly filled
 - Areas in Great Salt Lake, Utah
- Void Filling: (clockwise from upper left)
 - Original DEM
 - 2 km buffer around incorrect area
 - Extraction of cells to fill
 - Biharmonic Spline Interpolation



Common Datasets: Other Groups

- Invitation to Contribute through ~2020
 - Central America, South America, Caribbean Islands, W.
 Africa, Canary Islands, Samoa, Greenland, Iceland.

- Greenland (DTU) gravity data
- Iceland Landmælingar Íslands DEM/gravity data

geodesy.noaa.gov

Is it working?

GSVS11 – Texas	Minimum	Maximum	RMS	
[meters]				
xGEOID16A	-0.0381	0.0428	0.015	
xGEOID16B	-0.0365	0.0455	0.015	
xGEOID17A	-0.0494	0.0409	0.017	
xGEOID17B	-0.0344	0.0345	0.014	

- 44.0 2000 2000 0 0 0 0 0 0 0 0 0 0 0 0 0 0			ha w			xGEODTA xGEODTBB xGEODTA xGEODTA xGEODTA
4.04		1			11	VVV
0.04 0.04		100	Distance (br	4		xGEOID16A xGEOID16B xGEOID17A xGEOID17B
	MA	Antha	MANA	Munt	had	, AAA
4 11 - 4 12 -	V W	W	a where	ynd	Vinny	p wa
-0.00	V		V	- 4 C	M	
-3.05	30	100	150 Distance for	200	250	300

GSVS14 – Iowa	Minimum	Maximum	RMS
[meters]			
xGEOID16A	-0.031	0.053	0.015
xGEOID16B	-0.029	0.058	0.017
xGEOID17A	-0.025	0.035	0.013
xGEOID17B	-0.030	0.033	0.014

geodesy.noaa.gov

Contact Us:

Kevin M. Ahlgren, Ph.D. NGS Geoid Team

kevin.ahlgren@noaa.gov 240-533-9894

International Cooperation & Collaboration